



The Health Effects of Landslides

Introduction

Landslides are slope movements of rock, earth, mud, or debris that can deliver an array of devastating economic, social, and health impacts.¹ Landslides often accompany other disaster events, such as volcanic activity, flooding, earthquakes, tsunamis, or human activity such as construction, mining, or deforestation.² Such events can cause one singular landslide or trigger several landslides. Landslides may result in detrimental impacts to community health and well-being. Specifically, highly fluidized, rapid landslides with velocities greater than 5 m/s are typically the only kind of landslide Basic paragraph to result in detrimental human health impacts.³ Residents may be displaced from their homes and communities, need to seek emergency healthcare, and experience mental health conditions for months or years after the event. Additionally, landslides can cause unstable ground and create treacherous or impossible travel conditions, complicating evacuation and emergency response.^{4, 5}

Health Impacts

Acute and long-term landslide health impacts may have regional variation, depending on local culture, economics, and preparedness to disasters. Acute health impacts and mortality can occur during and immediately after landslides, both for at-risk and healthy individuals. Many affected by landslides experience a range of mental health impacts, which may not present for months following the event.⁶ While there has been limited research on health risks faced by recovery workers after a landslide event specifically, information on the physical and mental health risks experienced by disaster recovery workers more generally may inform precautions to protect their health and safety in this context.⁷⁻⁹

Acute Impacts

Flowing debris and mud can cause injuries during landslide events. Medical professionals characterize these injuries as “debris flow syndrome.”¹⁰ Other acute impacts may arise from displacement or harsher living conditions post-disaster, lack of clean water and sanitation, and airborne particulate matter in the lungs. Acute landslide impacts include:

- Soft tissue abrasions and lacerations, including to the eyes¹⁰
- Orthopedic injuries¹⁰
- Hypothermia¹⁰
- Craniofacial trauma¹⁰
- Mud impaction¹⁰
- Skin diseases¹¹
- Heat stroke¹¹
- Diarrhea¹¹
- Respiratory issues¹¹

Mental Health

Landslide survivors can experience mental health impacts for months to years after the event. In addition to physical injuries, many impacted individuals lose property, income, loved ones, or sense of place, and may experience economic challenges following the event.³ Individuals of all ages, including disaster response and recovery workers,¹² may be harmed psychologically by the event itself, even without other injury or loss.¹³ Some of the most common mental health impacts caused by landslides are:

- Post-traumatic stress disorder, or PTSD¹³
- Anxiety¹⁴
- Sleep disruption¹⁴
- Suicidal thoughts⁶

Mortality

Landslides and accompanying events have killed thousands of people globally.¹⁵ People who are driving, riding in trains, walking or otherwise outside, and in buildings that collapse are most likely to be fatally injured.¹⁶ Injuries that cause death include:

- Severe traumatic injuries¹⁷
- Suffocation¹⁸
- Complications from acute injuries¹⁸



Impacts to Healthcare

Because landslides can have a variety of impacts on different people, healthcare workers must be prepared to respond quickly and flexibly. Here are some lessons learned on the impacts on healthcare workers and healthcare infrastructure from past landslide events:

Impacts on Healthcare Workers

Communications infrastructure can be damaged during landslide events and cause delays in receiving outside aid.¹⁹

Healthcare workers may experience loss or injury during a landslide event, affecting their ability to support others.¹⁹

Patients may continue to seek medical care for disaster-related injuries days and weeks after the event.²⁰

Impacts on Healthcare

Some landslides have caused a surge in demand for medical assistance, in-patient care, emergency housing, and medical equipment, particularly in developing countries. Many healthcare facilities cannot handle such a surge due to lack of trained personnel, hospital bed capacity, storage to dispense controlled substances, and equipment such as CT scanners and ultrasounds.²¹

Landslides can impact roads and transportation systems, making it difficult to transport patients to healthcare facilities.²²



References

- Kennedy ITR, Petley DN, Williams R, Murray V. A systematic review of the health impacts of mass Earth movements (landslides). *PLoS Curr*. 2015;7:ecurrents.dis.1d49e84c8bbe678b0e70cf7fc35d0b77.
- Guzzetti F, Stark CP, Salvati P. Evaluation of flood and landslide risk to the population of Italy. *Environ Manage*. 2005;36(1):15-36.
- Pollock W, Wartman J. Human vulnerability to landslides. *GeoHealth*. 2020;4(10):e2020GH000287.
- Hungr O, Leroueil S, Picarelli L. The Varnes classification of landslide types, an update. *Landslides*. 2014;11(2):167-194.
- Hart A. Landslides. In: *Ciotton's Disaster Medicine*. Elsevier; 2024:640-643.
- Tang TC, Yen CF, Cheng CP, et al. Suicide risk and its correlate in adolescents who experienced typhoon-induced mudslides: a structural equation model. *Depress Anxiety*. 2010;27(12):1143-1148.
- Khatrri J, Fitzgerald G, Poudyal Chhetri MB. Health risks in disaster responders: A conceptual framework. *Prehosp Disaster Med*. 2019;34(2):209-216.
- Colf LA, McAleavy T. Health consequences of disasters: Advancing disaster data science. *PNAS Nexus*. 2024;3(6):gae211.
- Blackman D, Prayag G, Nakanishi H, Chaffer J, Freyens B. Wellbeing in disaster recovery: Understanding where systems get stuck. *Int J Disaster Risk Reduct*. 2023;95(103839):103839.
- Langdon S, Johnson A, Sharma R. Debris Flow Syndrome: Injuries and Outcomes after the Montecito Debris Flow. *Am Surg*. 2019;85(10):1094-1098.
- Chimed-Ochir O, Yumiya Y, Taji A, et al. Emergency Medical Teams' responses during the West Japan Heavy Rain 2018: J-SPEED data analysis. *Prehosp Disaster Med*. 2022;37(2):205-211.
- Mao X, Fung OWM, Hu X, Loke AY. Psychological impacts of disaster on rescue workers: A review of the literature. *Int J Disaster Risk Reduct*. 2018;27:602-617.
- Catapano F, Malafronte R, Lepre F, et al. Psychological consequences of the 1998 landslide in Sarno, Italy: a community study: Psychological consequences of Sarno landslide. *Acta Psychiatr Scand*. 2001;104(6):438-442.
- Moloney M, Quinkert E, PhD CH, PhD CB. 0114 "my mother doesn't sleep when it rains at night": Assessing the impacts of a climate-related flood on sleep. *Sleep*. 2024;47(Supplement_1):A50-A51.
- Petley D. Global patterns of loss of life from landslides. *Geology*. 2012;40(10):927-930.
- Salvati P, Petrucci O, Rossi M, Bianchi C, Pasqua AA, Guzzetti F. Gender, age and circumstances analysis of flood and landslide fatalities in Italy. *Sci Total Environ*. 2018;610-611:867-879.
- Uddin T, Islam MT, Gosney JE. 2017 Bangladesh landslides: physical rehabilitation perspective. *Disabil Rehabil*. 2021;43(5):718-725.
- Sanchez C, Lee TS, Young S, Batts D, Benjamin J, Malilay J. Risk factors for mortality during the 2002 landslides in Chuuk, Federated States of Micronesia. *Disasters*. 2009;33(4):705-720.
- Peltan ID. Disaster relief and recovery after a landslide at a small, rural hospital in Guatemala. *Prehosp Disaster Med*. 2009;24(6):542-548.
- Gathen M, Welle K, Jaenisch M, et al. Are orthopaedic surgeons prepared? An analysis of severe casualties from the 2021 flash flood and mudslide disaster in Germany. *European Journal of Trauma & Emergency Surgery*. 2022;48(5):4233-4241.
- Taylor-Robichaud A, Mugemangango P, Birahamire J, Martin KD. 370 emergency care capacity assessment of district hospitals in Rwanda using the WHO's hospital emergency assessment tool. *Ann Emerg Med*. 2024;84(4):S169.
- Pereira BMT, Morales W, Cardoso RG, Fiorelli R, Fraga GP, Briggs SM. Lessons learned from a landslide catastrophe in Rio de Janeiro, Brazil. *Am J Disaster Med*. 2013;8(4):253-258.

This material is based upon work supported by the National Science Foundation (NSF) under Awards #2103713 and #1940034. Additional support was provided by grant #P30ES007033 from the National Institute of Environmental Health Sciences (NIEHS). Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the NSF or NIEHS.

*Content by Naomi Cutler and Anna Kirchoff
Last updated 4/2025*